

Prenatal DDT Exposure in Relation to Anthropometric and Pubertal Measures in Adolescent Males

**Beth C. Gladen, Mark A. Klebanoff, Mary L. Hediger,
Solomon H. Katz, Dana B. Barr, Mark D. Davis,
and Matthew P. Longnecker**
doi:10.1289/ehp.7287 (available at <http://dx.doi.org/>)
Online 7 September 2004



Prenatal DDT Exposure in Relation to Anthropometric and Pubertal Measures in
Adolescent Males

Beth C. Gladen (1), Mark A. Klebanoff (2), Mary L. Hediger (2), Solomon H. Katz (3),
Dana B. Barr (4), Mark D. Davis (4), Matthew P. Longnecker (1)

(1) National Institute of Environmental Health Sciences, National Institutes of Health,
Department of Health and Human Services

(2) National Institute of Child Health and Human Development, National Institutes of
Health, Department of Health and Human Services

(3) Krogman Center for Research in Child Growth and Development, University of
Pennsylvania

(4) National Center for Environmental Health, Centers for Disease Control and
Prevention, Department of Health and Human Services

correspondence to: Beth C. Gladen

mailing address: Biostatistics Branch, Mail Drop A3-03

National Institute of Environmental Health Sciences, P.O. Box 12233

Research Triangle Park, NC 27709, USA

package delivery: Room A352, David P. Rall Building, 111 T.W. Alexander Drive

phone: 919-541-3461

fax: 919-541-4311

e-mail: gladen@niehs.nih.gov

running title: Prenatal DDT and Anthropometry and Puberty in Boys

key words:

DDT

DDE

Growth

Puberty

Child development

Prenatal exposure delayed effects

Acknowledgements:

A conflict of interest was not reported.

abbreviations:

BMI: body mass index

CPP: Collaborative Perinatal Project

DHEAS: Dehydroepiandrosterone sulfate

ND: not detected

o,p'-DDT: 1,1,1-trichloro-2-(*o*-chlorophenyl)-2-(*p*-chlorophenyl)-ethane

p,p'-DDE: 1,1-dichloro-2,2-bis(*p*-chlorophenyl)-ethylene

p,p'-DDT: 1,1,1-trichloro-2,2-bis(*p*-chlorophenyl)-ethane

PBPP: Philadelphia Blood Pressure Project

SEI: socioeconomic index

Section Headers:

Abstract

Introduction

Materials and Methods

Results

Discussion

References

Table 1

Table 2

Table 3

Table 4

Table 5

Abstract

DDT, a pesticide once used widely in agriculture and now limited to public health use, remains a controversial chemical due to a combination of benefits and risks. DDT or its breakdown products are ubiquitous in the environment and in humans. Compounds in the DDT family have endocrine actions and have been associated with reproductive toxicity. A previous study reported associations between prenatal exposure to p,p'-DDE and increased height and weight in adolescent boys. We examined a group with higher exposures to see whether similar associations would occur. Our study group was 304 males born in Philadelphia in the early 1960s who had participated in a previous study. Anthropometric and pubertal measures from one to six visits during their adolescent years were available, as were stored maternal serum samples from pregnancy. We measured p,p'-DDE, p,p'-DDT and o,p'-DDT in the maternal serum. Outcomes examined in the boys were height, ratio of sitting height to height, body mass index, triceps skinfold thickness, ratio of subscapular to the sum of triceps and subscapular skinfold thicknesses, skeletal age, serum testosterone, and serum dehydroepiandrosterone sulfate. No associations between prenatal exposure to any of the DDT compounds and any outcome measure were seen.

Introduction

The combination of public health benefits and environmental risks associated with DDT has made use of the pesticide controversial (Walker et al. 2003). DDT was once used extensively throughout the world, for both agricultural and public health purposes (ATSDR 2002). Usage today is generally limited to vector control, primarily of malaria, and is covered as of May 2004 by the Stockholm Convention on Persistent Organic Pollutants, a treaty signed by 151 countries and currently ratified by more than 75 countries (Stockholm Convention on Persistent Organic Pollutants, 2004). Decisions in any specific instance about whether to use DDT even for vector control can be hotly debated (Wendo 2004).

Usage of DDT in the USA began in the 1940s, peaked in the early 1960s, and essentially ended in 1972. Timelines in other countries differed, but generally followed the same pattern of steep rise and fall in amounts used. Despite the decline in use, several components and breakdown products of the pesticide are still widely detectable in the environment and in humans (Jaga and Dharmani 2003; Smith 1999). The pesticide product consists primarily of the actual insecticide p,p'-DDT, with o,p'-DDT and several other minor components making up the remainder (ATSDR 2002). The primary degradation product and human metabolite of p,p'-DDT is p,p'-DDE; the latter is also the most persistent of the DDT family and the one that bioaccumulates most extensively in humans.

The various components of the DDT family have a number of known biological actions. The insecticidal effects of p,p'-DDT are attributable to neurotoxicity. A number of endocrine effects have been shown, including antiandrogenic properties of p,p'-DDE (Gray et al. 2001), estrogenic properties of o,p'-DDT (Kupfer 1975), and modulation of steroid hormone homeostasis through induction of hepatic enzymes (Wyde et al. 2003; You et al. 2001). Changes

in immune markers have been seen (Vine et al. 2001). Carcinogenicity has been demonstrated in animals, although evidence in humans is mixed (Turusov et al. 2002). Associations with impaired lactation have been reported (Gladen and Rogan 1995; Rogan et al. 1987). Reproductive effects have been shown. A large human study has shown associations of maternal p,p'-DDE with preterm birth and decreased birth weight (Longnecker et al. 2001). Animal studies with p,p'-DDE show a number of reproductive abnormalities in male offspring (Gray et al. 2001); whether similar effects are seen in humans at usual exposure levels is uncertain (Flores-Luévano et al. 2003; Hosie et al. 2000; Longnecker et al. 2002). Paternal occupational DDT exposure has also been associated with birth defects (Salazar-García et al. 2004).

Prenatal exposure to DDT can also have delayed effects. There are reports of neurotoxic and immunotoxic effects in young children (Dewailly et al. 2000; Ribas-Fitó et al. 2003), although not all studies show such effects (Gladen et al. 1988; Rogan et al. 1987). At even later ages, prenatal and lactational exposure to p,p'-DDE in animals has been associated with delayed male puberty in some but not all studies (Loeffler and Peterson 1999; You et al. 1998).

A previous human study had shown that adolescent males with higher prenatal exposure to p,p'-DDE had increases in both height and body mass index compared to those with lower exposures; markers of puberty were unaffected (Gladen et al. 2000). In view of the continuing controversy about DDT and concerns about substantial increases in childhood obesity (Ogden et al. 2002), we examined another population with higher exposures to see whether similar effects in adolescent males would be seen there.

Materials and Methods

Subjects

The Collaborative Perinatal Project (CPP) was a large multi-center prospective study of approximately 50,000 children born between 1959 and 1966 (Broman 1984; Hardy 2003). Data collected included background information on the mothers obtained from questionnaires during pregnancy. Serum samples taken from the mothers during pregnancy were stored, and many are still available.

The Philadelphia Blood Pressure Project (PBPP) followed some of the children enrolled at one of the CPP centers during adolescence and early adulthood (Katz et al. 1980). This was an urban population; the children had been born at Pennsylvania Hospital and followed at Children's Hospital of Philadelphia. There were several subgroups chosen for study; one was a stratified random sample of those enrolled in the CPP who were born between 1961 and 1965. These subjects were seen annually up to 3 times in 1977-80 and again annually up to another 3 times in 1982-85. Data collected included anthropometric measurements and pubertal markers.

The subjects of the present study were chosen from among the 373 singleton males from the random sample studied in the PBPP. Of those, 314 had stored maternal samples from the third trimester available. Those samples were shipped to the Centers for Disease Control and Prevention and analyzed for several DDT compounds. Of the samples shipped, 9 had insufficient quantity for analysis, 1 was lost during analysis, and 304 were successfully analyzed. The 304 boys whose maternal samples were analyzed are the subjects of this report.

Primary Variables

Anthropometric measurements up to age 20 years from the PBPP were used; data from a total of 1137 visits from the 304 boys were available. Measurement techniques and reliabilities

have been discussed previously (Katz et al. 1980; Tanner et al. 1969). Quantities measured at all visits included height, sitting height, weight, triceps skinfold thickness, and subscapular skinfold thickness. Height and sitting height were measured using a Holtain stadiometer. Weight was measured on a Homs beam scale. Skinfolts were measured using a Holtain skinfold caliper. We examined several measures of overall size and body proportion; these included height, the ratio of sitting height to height (height ratio), body mass index (BMI), triceps skinfold thickness, and the ratio of subscapular to the sum of subscapular and triceps skinfold thicknesses (a measure of central adiposity). Missing data were minimal; height was unavailable 4 times, sitting height 14 times, weight 7 times, triceps skinfold thickness 4 times, and subscapular skinfold thickness 7 times.

Skeletal age was determined at the three PBPP visits in 1977-80, using the Tanner-Whitehouse II method of rating hand-wrist radiographs on the maturity of 20 individual bones (Katz et al. 1980; Tanner et al. 1975). Skeletal age was unavailable for 4% of the visits where it was scheduled to be done.

Testosterone was measured at the first two PBPP visits in 1977-79 by radioimmunoassay on samples of venous blood collected at the time of examination (Furuyama et al. 1970; Zemel and Katz 1986). Dehydroepiandrosterone sulfate (DHEAS) was measured only at the second visit in 1978-79, again by radioimmunoassay (Buster and Abraham 1972). Not all boys volunteered for the blood draw; testosterone was unavailable for 23% of the visits where it was scheduled to be done, and DHEAS was unavailable in 22%.

Maternal serum samples were analyzed for p,p'-DDE (1,1-dichloro-2,2-bis(*p*-chlorophenyl)-ethylene), p,p'-DDT (1,1,1-trichloro-2,2-bis(*p*-chlorophenyl)-ethane), and o,p'-DDT (1,1,1-trichloro-2-(*o*-chlorophenyl)-2-(*p*-chlorophenyl)-ethane) using a semi-automated

solid phase extraction and gel permeation chromatography cleanup followed by an isotope dilution gas chromatography-high resolution mass spectrometry analysis (Barr et al. 2003; Sandau et al. 2003). Recovery correction was done for each analyte in each individual sample. For 25 samples, p,p'-DDT could not be measured due to quality control limit failure. Cholesterol and triglycerides were also measured using standard clinical assays. Total serum lipids were calculated as $62.3 + 2.27 \text{ cholesterol} + \text{triglycerides}$ (Phillips et al. 1989). Pesticide concentrations were reported as ng pesticide per g total serum lipids. The sum of the three DDT compounds was calculated, unless p,p'-DDT could not be measured. Samples with non-detectable amounts for o,p'-DDT or p,p'-DDT were considered to be zero; since detection limits were low, imputing any other value up to the detection limit changed the sum of DDT by less than 1% and never changed the categories when exposure was discretized.

Statistical Analysis

Models were used to examine outcome measurements in relation to pesticide concentration after adjustment for important predictors and potential confounders. The age of the boy at examination is a key predictor and was included as a cubic polynomial to allow for nonlinearity. Parental size is also a strong predictor of child size; maternal height and prepregnancy BMI were available and were included as linear terms, but paternal size was not available. We also adjusted for breast feeding (yes, no), maternal smoking at the time of pregnancy (yes, no), number of older siblings (0, 1, 2+), race (black, white), maternal age at birth (13-19, 20-24, 25-29, 30+ years), and maternal age at menarche (8-11, 12, 13, 14+). We used the family socioeconomic index (SEI) score, based on education, occupation, and income, created by the CPP investigators for internal comparisons (Myrianthopoulos and French 1968). Median SEI for the entire CPP was 4.3, with a range from 0 to 9.5; scores here were categorized into three

groups (0-2.5, 2.6-5.0, 5.1+). A term for each boy was included as a random effect to account for the correlation among multiple measures of the same boy. For triceps skinfold thickness and testosterone, the analysis was done on a log scale. Tests of statistical significance reported are either tests of whether categories differ for discrete predictors or tests for zero slope for continuous predictors. Models were fit using SAS version 9 (SAS Institute, Cary NC).

Results

Most of the 304 boys in this study were black (Table 1). A majority of their mothers reported smoking during pregnancy and few mothers breast-fed their sons. A quarter of the boys were first-born. The distribution of family SEI was similar to that of all blacks in the CPP. Other characteristics of the mother and family around the time of birth are shown in Table 1.

Concentrations of p,p'-DDE in maternal serum during pregnancy ranged from 1 to 25 $\mu\text{g/g}$ lipid (Table 2), with a median of 5.7 $\mu\text{g/g}$ lipid. The other two DDT compounds were present at lower concentrations; median p,p'-DDT was 1.9 $\mu\text{g/g}$ lipid and median o,p'-DDT was 0.14 $\mu\text{g/g}$ lipid. The three compounds measured were correlated; the correlation of p,p'-DDE with p,p'-DDT was 0.65 and with o,p'-DDT was 0.58, while p,p'-DDT and o,p'-DDT had a correlation of 0.77.

The boys had from one to six adolescent visits with anthropometric measurements available; 20% had one or two, 34% had three, and 47% had four or more. Age at the first measurement ranged from 10.8 to 17.9 years, with a median of 12.8 years. Age at the last measurement ranged from 12.2 to 20.0 years, with a median of 17.6 years. Age at first measurement was a major determinant of number of measurements, as those who were older at the start of followup left the age range of interest more quickly. In addition, no whites had more than three visits.

Height ranged from 132 to 196 cm, with the expected strong relationship to age. Selected percentiles at each age are shown in Table 3. Height ratio ranged from 46 to 55%. BMI was skewed, ranging from 14 to 45 kg/m². Triceps skinfold was more skewed, ranging from 3.5 to 43.4 mm. Central adiposity ranged from 31 to 78%.

The boys also had up to three measures of skeletal age; 66% had all three measurements, and two boys had none. Skeletal age ranged from 8.7 to 18 years. In addition, 260 boys had one or two testosterone measurements available and 213 had a single measurement of DHEAS. Testosterone ranged from 1 to 1258 ng/dl; DHEAS ranged from 56 to 5600 ng/ml.

The crude relationships of these anthropometric and pubertal measures to maternal prenatal concentrations of p,p'-DDE are shown in table 4 for several age ranges. Not all boys had measurements available in all age ranges; for skeletal ages and hormones, there were few measurements past age 17. Little systematic relationship to p,p'-DDE was seen for any of the measures.

Models as described above were fit to these measures to allow age to be treated as a continuous predictor and to adjust for other predictors and potential confounders (table 5). In no case was p,p'-DDE a statistically significant predictor of the outcome (all $p > 0.10$). Effects of other predictors were seen. All outcomes were significantly related to the age of the boy at measurement (data not shown). Height also increased with maternal height ($p < 0.001$) and maternal BMI ($p = 0.053$); first born boys also had greater means ($p = 0.043$), as did those from families with higher SEI ($p = 0.094$). Height ratio decreased with maternal height ($p < 0.001$); whites also had larger means ($p < 0.001$), as did later-born children ($p = 0.063$) and those whose mothers had early menarche ($p = 0.095$). BMI increased with maternal BMI ($p < 0.001$). Triceps skinfold increased with maternal BMI ($p < 0.001$); first born boys ($p = 0.037$) and whites ($p = 0.056$)

had higher means. Mean central adiposity was higher in blacks ($p=0.002$). Testosterone was increased among those whose mothers had early menarche ($p=0.052$) and those from families with higher SEI ($p=0.092$). Skeletal age and DHEAS showed no significant effects of predictors other than age of the boy.

Use of p,p'-DDT, o,p'-DDT, or the sum of the three compounds rather than p,p'-DDE as the exposure also resulted in no significant effects on any of the outcomes analyzed (data not shown). When the analysis shown in table 5 was done separately for each of the age groups used in table 4, the results were again not significant with one exception. At the youngest ages, the 5 exposure groups had significantly different BMIs, but the pattern was not monotonic in dose; as with the crude results in table 4, the highest BMIs were seen for the 3-6 dose group. If the analysis shown in table 5 is restricted to blacks, the results are essentially unchanged (data not shown).

Discussion

In this study, we found no association of prenatal exposure to p,p'-DDE, p,p'-DDT, or o,p'-DDT with any of the anthropometric or pubertal measures we examined in adolescent males. In a previous study of 278 adolescent boys and 316 girls, prenatal p,p'-DDE exposure was also not related to pubertal markers (Gladen et al. 2000). However, increased exposure in that study was associated with greater height and BMI of the boys. The subjects of the present study, who were born during the peak of DDT usage in the USA, had higher exposures than those in the previous study, who were born after agricultural DDT use had been banned. Median p,p'-DDE in maternal serum in the current study was 5.7 $\mu\text{g/g}$ serum lipid; the median in the previous study was approximately equivalent to 1.6 $\mu\text{g/g}$ serum lipid (12.6 ng/g serum (Rogan et al. 1986), converted assuming 8 g lipid/l serum (Longnecker et al. 2003)). The failure to replicate the

previous findings on height and BMI in the present study with higher exposures raises the possibility that the earlier results may have been due to chance. However, we note that there were a number of differences in the populations studied; for example, the previous study subjects were mostly whites, mostly breast-fed, and had mothers who were of higher socioeconomic status and less likely to smoke.

Childhood concentration of p,p'-DDE has also been studied in relation to childhood height and pubertal development (Denham et al. 2004; Karmaus et al. 2002). Adult concentration of p,p'-DDE has also been studied in relation to testosterone and DHEAS (Ayotte et al. 2001; Hagmar et al. 2001; Martin et al. 2002; Persky et al. 2001). However, these studies have limited relevance to the question addressed here. Prenatal exposure is likely to act through different mechanisms than does postnatal exposure. Childhood concentrations of persistent organochlorines like the DDT compounds are poor surrogates for prenatal exposures, as concentrations even into adolescence are most strongly determined by breast-feeding (Jacobson et al. 1989; Karmaus et al. 2001; Nawrot et al. 2002).

The adolescent period studied here is a time of rapid development, with changes in body size and proportions, development of secondary sexual characteristics, skeletal maturation, and changes in the hormonal milieu all occurring. Prenatal exposure to compounds with endocrine activity might influence either the timing or the ultimate result of any or all of these changes. All of the outcome measures we studied reflect some aspect of adolescent development, although we do lack some classic outcomes such as Tanner stages and time of peak height velocity. Height increases with age, although it levels off in the later teens; height ratio first declines and then increases with age as body proportions shift (Hamill et al. 1973; Malina et al. 1974). BMI increases with age, albeit with considerable variability. Triceps skinfold thickness declines with

age, again with considerable variability; it also becomes smaller relative to subscapular skinfold thickness, so that central adiposity increases. Skeletal age increases with chronological age, up to full maturity at skeletal age 18. Testosterone and DHEAS concentrations increase with age, with considerable variability. The substantial variability seen with some of these measures means that our failure to find associations of prenatal exposure to the DDT compounds with any of these outcomes could be due to inadequate power, although the patterns of the observed relationships do not suggest this explanation.

We did have adequate power to discern effects of other known predictors. As expected, height and BMI of boys was influenced by the height and BMI of their mothers, consistent with previous work (Celi et al. 2003; Wingerd and Schoen 1974). The racial differences we saw were consistent with those seen elsewhere. In a national survey, the relationship of height and weight to race was inconsistent across age, but height ratio showed a clear racial difference (Hamill et al. 1973). In the same survey, whites had greater triceps skinfolds than blacks but similar subscapular skinfolds, leading to lower central adiposity (Johnston et al. 1974); racial differences in skeletal age were inconsistent across ages (Roche et al. 1975, 1978). First-born children have been shown to be taller and heavier (Celi et al. 2003; Ong et al. 2002; Wingerd and Schoen 1974), consistent with our findings; increased skinfold thickness among first-borns has also been reported in another investigation based on the PBPP (Stettler et al. 2000). Prenatal exposure to smoking has been associated with decreased height (Fogelman 1980) and increased obesity (Power and Jefferis 2002); our results were in the expected direction, although they did not achieve statistical significance. Our sample included very few breast-fed children, consistent with the low overall breast-feeding rates at that time and with the lower rates in blacks and in the

northeast (Hirschman and Hendershot 1979), so the failure to see any associations of our outcomes to breast-feeding was not surprising.

While there are many early influences on later development, such as the association of prenatal exposure to certain anti-psychotic drugs with later height (Platt et al. 1988), most would be expected to be unrelated to exposure to DDT and thus not candidate confounders. We controlled for the most likely confounders, although observational studies are always subject to potential residual confounding. We had no information about maternal diet prior to pregnancy nor about the diet of the child after birth; maternal diet is a predictor of the exposure and childhood diet is a predictor of growth and development. Adjusting for paternal size might have made our estimates more precise, but this information was not available. Birth weight has been shown to be related to prenatal p,p'-DDE exposure in the CPP population (Longnecker et al. 2001); however, we did not consider it appropriate to adjust for birth weight since this is an intermediate variable in the relationships between exposure and adolescent outcomes.

Exposure was measured using third-trimester serum samples. Specific aspects of prenatal development occur during critical windows, so timing of exposure can be important. However, for persistent compounds like DDT, concentrations are generally stable over periods of months or longer, with little variation over the course of pregnancy (Longnecker et al. 1999). The analytical methods used were sensitive, selective, and reliable, with relative standard deviations, including both the error from the sample preparation and the instrumental methods, of 11% and detection limits in the low pg/ml range (Barr et al. 2003).

The participants in the study were not a random sample of the general population. Those enrolled in the CPP from the Philadelphia study center were clinic patients who were planning to deliver at the study hospital (Broman 1984); they were mostly black and relatively low income,

representative of the population obtaining medical care at this clinic. The PBPP study group was a random sample of the CPP study group (Katz et al. 1980). We have no information about exposure among those who chose not to participate in either the base CPP study or the PBPP followup, but there is no reason to anticipate that prenatal DDT exposure would differ between participants and non-participants. Refusal rates were greatest for the hormone measurements, but there was little systematic relationship to exposure; for example, among blacks, those with both testosterone measurements available had a median p,p'-DDE of 6.2, while those with none or one, due to either a missed visit or a refusal, had a median of 6.3.

In summary, we have seen no association between prenatal exposure to DDT-related compounds and several anthropometric and pubertal measures in males. However, high variability in some of the outcome measures means we cannot rule out subtle changes.

References

- ATSDR. 2002. Toxicological Profile for DDT, DDE, and DDD. Atlanta, GA: Agency for Toxic Substances and Disease Registry.
- Ayotte P, Giroux S, Dewailly É, Hernández Avila M, Farias P, Danis R, et al. 2001. DDT spraying for malaria control and reproductive function in Mexican men. *Epidemiology* 12:366-367.
- Barr JR, Maggio VL, Barr DB, Turner WE, Sjödin A, Sandau CD, et al. 2003. New high-resolution mass spectrometric approach for the measurement of polychlorinated biphenyls and organochlorine pesticides in human serum. *J Chromatogr B Analyt Technol Biomed Life Sci* 794:137-148.
- Broman S. 1984. The Collaborative Perinatal Project: An overview. In: *Handbook of Longitudinal Research* (Medrick SA, Harway M, Finello KM, eds). New York: Praeger Publications, 185-215.
- Buster JE, Abraham GE. 1972. Radioimmunoassay of plasma dehydroepiandrosterone sulfate. *Anal Lett* 5:543-551.
- Celi F, Bini V, DeGiorgi G, Molinari D, Faraoni F, DiStefano G, et al. 2003. Epidemiology of overweight and obesity among school children and adolescents in three provinces of central Italy, 1993–2001: study of potential influencing variables. *Eur J Clin Nutr* 57:1045-1051.
- Denham M, Deane G, Gallo MV, Ravenscroft J, Decaprio A, Schell L. 2004. Age at menarche among Akwesasne Mohawk girls exposed to lead, mercury, mirex, dichlorodiphenyldichloroethane (DDE), hexachlorobenzene (HCB) and polychlorinated biphenyls (PCBs) [Abstract]. *Am J Hum Biol* 16:201.

- Dewailly É, Ayotte P, Bruneau S, Gingras S, Belles-Isles M, Roy R. 2000. Susceptibility to infections and immune status in Inuit infants exposed to organochlorines. *Environ Health Perspect* 108:205-211.
- Flores-Luévano S, Farías P, Hernández M, Romano-Riquer P, Weber JP, Dewailly E, et al. 2003. Concentraciones de DDT/DDE y riesgo de hipospadias. Un estudio piloto de casos y controles [in Spanish]. *Salud Publica Mex* 45:431-438.
- Fogelman K. 1980. Smoking in pregnancy and subsequent development of the child. *Child Care Health Dev* 6:233-249.
- Furuyama S, Mayes DM, Nugent CA. 1970. A radioimmunoassay for plasma testosterone. *Steroids* 16:415-428.
- Gladen BC, Ragan NB, Rogan WJ. 2000. Pubertal growth and development and prenatal and lactational exposure to PCBs and DDE. *J Pediatr* 136:490-496.
- Gladen BC, Rogan WJ. 1995. DDE and shortened duration of lactation in a northern Mexican town. *Am J Public Health* 85:504-508.
- Gladen BC, Rogan WJ, Hardy P, Thullen J, Tingelstad J, Tully M. 1988. Development after exposure to polychlorinated biphenyls and dichlorodiphenyl dichloroethene transplacentally and through human milk. *J Pediatr* 113:991-995.
- Gray LE, Ostby J, Furr J, Wolf CJ, Lambright C, Parks L, et al. 2001. Effects of environmental antiandrogens on reproductive development in experimental animals. *Hum Reprod Update* 7:248-264.
- Hagmar L, Björk J, Sjödin A, Bergman Å, Erfurth EM. 2001. Plasma levels of persistent organohalogenes and hormone levels in adult male humans. *Arch Environ Health* 56:138-143.

- Hamill PVV, Johnston FE, Lemeshow S. 1973. Body weight, stature, and sitting height: White and Negro youths 12-17 years: United States. *Vital Health Stat* 11 126:1-40.
- Hardy JB. 2003. The Collaborative Perinatal Project: Lessons and legacy. *Ann Epidemiol* 13:303-311.
- Hirschman C, Hendershot GE. 1979. Trends in Breast Feeding among American Mothers. *Vital and Health Statistics, Series 23* 3:1-46.
- Hosie S, Loff S, Witt K, Niessen K, Waag KL. 2000. Is there a correlation between organochlorine compounds and undescended testes? *Eur J Pediatr Surg* 10:304-309.
- Jacobson JL, Humphrey HEB, Jacobson SW, Schantz SL, Mullin MD, Welch R. 1989. Determinants of polychlorinated biphenyls (PCBs), polybrominated biphenyls (PBBs), and dichlorodiphenyl trichloroethane (DDT) levels in the sera of young children. *Am J Public Health* 79:1401-1404.
- Jaga K, Dharmani C. 2003. Global surveillance of DDT and DDE levels in human tissues. *Int J Occup Med Environ Health* 16:7-20.
- Johnston EE, Hamill PVV, Lemeshow S. 1974. Skinfold thickness of youths 12-17 years, United States. *Vital Health Stat* 11 132:1-75.
- Karmaus W, Asakevich S, Indurkha A, Witten J, Kruse H. 2002. Childhood growth and exposure to dichlorodiphenyl dichloroethene and polychlorinated biphenyls. *J Pediatr* 140:33-39.
- Karmaus W, DeKoning EP, Kruse H, Witten J, Osius N. 2001. Early childhood determinants of organochlorine concentrations in school-aged children. *Pediatr Res* 50:331-336.

- Katz SH, Hediger ML, Schall JJ, Bowers EJ, Barker WF, Aurand S, et al. 1980. Blood pressure, growth and maturation from childhood through adolescence: Mixed longitudinal analyses of the Philadelphia Blood Pressure Project. *Hypertension* 2:I55-I69.
- Kupfer D. 1975. Effects of pesticides and related compounds on steroid metabolism and function. *Crit Rev Toxicol* 4:83-124.
- Loeffler IK, Peterson RE. 1999. Interactive effects of TCDD and p,p'-DDE on male reproductive tract development in in utero and lactationally exposed rats. *Toxicol Appl Pharmacol* 154:28-39.
- Longnecker MP, Klebanoff MA, Brock JW, Zhou H, Gray KA, Needham LL, et al. 2002. Maternal serum level of 1,1-dichloro-2,2-bis(p-chlorophenyl)ethylene and risk of cryptorchidism, hypospadias, and polythelia among male offspring. *Am J Epidemiol* 155:313-322.
- Longnecker MP, Klebanoff MA, Gladen BC, Berendes HW. 1999. Serial levels of serum organochlorines during pregnancy and postpartum. *Arch Environ Health* 54:110-114.
- Longnecker MP, Klebanoff MA, Zhou H, Brock JW. 2001. Association between maternal serum concentration of the DDT metabolite DDE and preterm and small-for-gestational-age babies at birth. *Lancet* 358:110-114.
- Longnecker MP, Wolff MS, Gladen BC, Brock JW, Grandjean P, Jacobson JL, et al. 2003. Comparison of polychlorinated biphenyl levels across studies of human neurodevelopment. *Environ Health Perspect* 111:65-70.
- Malina RM, Hamill PVV, Lemeshow S. 1974. Body dimensions and proportions, white and negro children, 6-11 years, United States. *Vital Health Stat* 11 143:1-71.

- Martin SA, Jr., Harlow SD, Sowers MF, Longnecker MP, Garabrant D, Shore DL, et al. 2002. DDT metabolite and androgens in African-American farmers. *Epidemiology* 13:454-458.
- Myrianthopoulos NC, French KS. 1968. An application of the U.S. Bureau of the Census socioeconomic index to a large, diversified patient population. *Soc Sci Med* 2:283-299.
- Nawrot TS, Staessen JA, Den Hond EM, Koppen G, Schoeters G, Fagard R, et al. 2002. Host and environmental determinants of polychlorinated aromatic hydrocarbons in serum of adolescents. *Environ Health Perspect* 110:583-589.
- Ogden CL, Flegal KM, Carroll MD, Johnson CL. 2002. Prevalence and trends in overweight among US children and adolescents, 1999-2000. *JAMA* 288:1728-1732.
- Ong KKL, Preece MA, Emmett PM, Ahmed ML, Dunger DB, ALSPAC Study Team. 2002. Size at birth and early childhood growth in relation to maternal smoking, parity and infant breast-feeding: longitudinal birth cohort study and analysis. *Pediatr Res* 52:863-867.
- Persky V, Turyk M, Anderson HA, Hanrahan LP, Falk C, Steenport DN, et al. 2001. The effects of PCB exposure and fish consumption on endogenous hormones. *Environ Health Perspect* 109:1275-1283.
- Phillips DL, Pirkle JL, Burse VW, Bernert JT, Jr, Henderson LO, Needham LL. 1989. Chlorinated hydrocarbon levels in human serum: effects of fasting and feeding. *Arch Environ Contam Toxicol* 18:495-500.
- Platt JE, Friedhoff AJ, Broman SH, Bond RN, Laska E, Lin SP. 1988. Effects of prenatal exposure to neuroleptic drugs on children's growth. *Neuropsychopharmacology* 1:205-212.
- Power C, Jefferis BJ. 2002. Fetal environment and subsequent obesity: a study of maternal smoking. *Int J Epidemiol* 31:413-419.

- Ribas-Fitó N, Cardo E, Sala M, de Muga ME, Mazón C, Verdú A, et al. 2003. Breastfeeding, exposure to organochlorine compounds, and neurodevelopment in infants. *Pediatrics* 111:e580-e585.
- Roche AF, Roberts J, Hamill PVV. 1975. Skeletal maturity of children 6-11 years: Racial, geographic area, and socioeconomic differentials: United States. *Vital Health Stat* 11 149:1-88.
- — —. 1978. Skeletal maturity of youths 12-17 years: Racial, geographic area, and socioeconomic differentials: United States, 1966-1970. *Vital Health Stat* 11 167:1-107.
- Rogan WJ, Gladen BC, McKinney JD, Carreras N, Hardy P, Thullen J, et al. 1986. Polychlorinated biphenyls (PCBs) and dichlorodiphenyl dichloroethene (DDE) in human milk: Effects of maternal factors and previous lactation. *Am J Public Health* 76:172-177.
- — —. 1987. Polychlorinated biphenyls (PCBs) and dichlorodiphenyl dichloroethene (DDE) in human milk: Effects on growth, morbidity, and duration of lactation. *Am J Public Health* 77:1294-1297.
- Salazar-García F, Gallardo-Díaz E, Cerón-Mireles P, Loomis D, Borja-Aburto VH. 2004. Reproductive effects of occupational DDT exposure among male malaria control workers. *Environ Health Perspect* 112:542-547.
- Sandau CD, Sjödin A, Davis MD, Barr JR, Maggio VL, Waterman AL, et al. 2003. Comprehensive solid-phase extraction method for persistent organic pollutants. Validation and application to the analysis of persistent chlorinated pesticides. *Anal Chem* 75:71-77.
- Smith D. 1999. Worldwide trends in DDT levels in human breast milk. *Int J Epidemiol* 28:179-188.

- Stettler N, Tershakovec AM, Zemel BS, Leonard MB, Boston RC, Katz SH, et al. 2000. Early risk factors for increased adiposity: a cohort study of African American subjects followed from birth to young adulthood. *Am J Clin Nutr* 72:378-383.
- Stockholm Convention on Persistent Organic Pollutants. 2004. Available: <http://www.pops.int> [accessed 24 August 2004]
- Tanner JM, Hiernaux J, Jarman S. 1969. Growth and physique studies. In: *Human Biology, A Guide to Field Methods*, IBP Handbook No. 9 (Weiner JS, Lourie JA, eds). Oxford: Blackwell, International Biological Programme.
- Tanner JM, Whitehouse RH, Marshall WA, Healy MJR, Goldstein H. 1975. Assessment of skeletal maturity and prediction of adult height (Tanner-Whitehouse II method). London: Academic Press.
- Turusov V, Rakitsky V, Tomatis L. 2002. Dichlorodiphenyltrichloroethane (DDT): Ubiquity, persistence, and risks. *Environ Health Perspect* 110:125-128.
- Vine MF, Stein L, Weigle K, Schroeder J, Degnan D, Tse CK, et al. 2001. Plasma 1,1-dichloro-2,2-bis(p-chlorophenyl)ethylene (DDE) levels and immune response. *Am J Epidemiol* 153:53-63.
- Walker KR, Ricciardone MD, Jensen J. 2003. Developing an international consensus on DDT: a balance of environmental protection and disease control. *Int J Hyg Environ Health* 206:423-435.
- Wendo C. 2004. Uganda considers DDT to protect homes from malaria. Health officials claim DDT will help save money, but critics warn of environmental costs. *Lancet* 363:1376.
- Wingard J, Schoen EJ. 1974. Factors influencing length at birth and height at five years. *Pediatrics* 53:737-741.

- Wyde ME, Bartolucci E, Ueda A, Zhang H, Yan B, Negishi M, et al. 2003. The environmental pollutant 1,1-dichloro-2,2-bis (p-chlorophenyl)ethylene induces rat hepatic cytochrome P450 2B and 3A expression through the constitutive androstane receptor and pregnane X receptor. *Mol Pharmacol* 64:474-481.
- You L, Casanova M, Archibeque-Engle S, Sar M, Fan LQ, Heck HD. 1998. Impaired male sexual development in perinatal Sprague-Dawley and Long-Evans Hooded rats exposed *in utero* and lactationally to *p,p'*-DDE. *Toxicol Sci* 45:162-173.
- You L, Sar M, Bartolucci E, Ploch S, Whitt M. 2001. Induction of hepatic aromatase by *p,p'*-DDE in adult male rats. *Mol Cell Endocrinol* 178:207-214.
- Zemel BS, Katz SH. 1986. The contribution of adrenal and gonadal androgens to the growth in height of adolescent males. *Am J Phys Anthropol* 71:459-466.

Table 1: Background characteristics of boys and their families

Characteristic	Value	Percent
Race	white	15
	black	85
Maternal smoking at time of pregnancy	no	44
	yes	56
Breast-fed	yes	6
	no	94
Number of older siblings	0	25
	1	25
	2+	50
Maternal height (cm)	144-<156	29
	156-<166	54
	166- 181	17
Maternal prepregnancy BMI (kg/m ²)	16-<20	21
	20-<25	47
	25-<30	23
	30- 44	9
Maternal age at menarche (yrs)	8-11	21
	12	29
	13	25
	14+	26
Maternal age at enrollment in CPP (yrs)	13-19	23
	20-24	36
	25-29	23
	30-42	18
Family SEI at time of pregnancy	0-2.5	17
	2.6-5.0	59
	5.1+	24

Number of cases (out of 304) with missing data: 3 for breast-fed, 1 for number of older siblings, 4 for maternal height, 6 for maternal prepregnancy BMI, 2 for maternal age at menarche, 10 for family SEI at time of pregnancy.

Table 2: Distribution of chemical concentrations in maternal serum

Chemical	Concentration, $\mu\text{g/g}$ lipid	Percent
p,p'-DDE	1.0- 2.9	14
	3.0- 5.9	38
	6.0- 8.9	24
	9.0-11.9	13
	12.0-25.1	11
p,p'-DDT ^a	ND ^b -0.9	17
	1.0- 1.9	36
	2.0- 2.9	24
	3.0- 3.9	11
	4.0-12.7	11
o,p'-DDT	ND ^c -0.07	29
	0.08-0.15	28
	0.16-0.23	18
	0.24-0.31	10
	0.32-1.33	14
sum DDT ^a	1.8- 3.9	11
	4.0- 7.9	37
	8.0-11.9	28
	12.0-15.9	13
	16.0-33.1	11

ND denotes not detected.

^a Not available for 25 (out of 304) boys due to quality control limit failure.

^b 1 sample had non-detectable p,p'-DDT (detection limit 0.01 $\mu\text{g/g}$ lipid).

^c 15 samples had non-detectable o,p'-DDT (detection limits 0.007 to 0.054 $\mu\text{g/g}$ lipid).

Table 3: Percentiles of anthropometric and pubertal measures by age

Age (yrs)	10-10.9	11-11.9	12-12.9	13-13.9	14-14.9	15-15.9	16-16.9	17-17.9	18-18.9	19-20.0
height (cm)										
N	10	97	149	185	140	111	105	118	111	107
90th	-	157	165	171	177	180	184	185	185	185
median	140	145	151	158	166	171	173	173	175	176
10th	-	138	143	147	154	158	166	167	168	167
height ratio (%)										
N	10	97	148	185	140	110	103	117	108	105
90th	-	52.9	52.5	52.3	52.2	53.0	52.6	52.7	52.4	52.3
median	52.0	51.1	50.5	50.3	50.1	50.2	50.5	50.8	50.8	50.8
10th	-	49.4	49.0	48.8	48.7	48.3	48.4	49.0	49.1	49.4
BMI (kg/m ²)										
N	10	97	149	185	139	111	105	118	110	106
90th	-	22.0	23.9	23.7	24.2	25.2	24.6	27.5	27.5	27.8
median	17.2	17.4	18.0	18.6	19.3	20.4	20.4	21.4	22.2	21.8
10th	-	15.2	15.4	15.9	16.1	17.6	17.9	19.0	19.2	19.1
triceps skinfold thickness (mm)										
N	10	97	149	185	140	111	105	118	111	107
90th	-	16.9	18.2	16.7	16.1	17.1	10.3	15.0	15.3	14.6
median	7.8	8.1	7.8	7.4	7.1	7.0	6.6	6.8	7.3	6.9
10th	-	5.6	5.2	4.9	5.0	5.2	4.6	5.0	4.9	4.5

central adiposity (%)										
N	9	97	148	185	140	110	105	118	111	107
90th	-	52	54	56	56	58	60	63	64	65
median	44	46	47	48	49	52	55	56	56	58
10th	-	40	41	41	42	42	46	49	49	50
skeletal age (years)										
N	9	95	146	177	135	106	78	37	4	0
90th	-	13.5	15.0	15.3	16.1	18.0	18.0	18.0	-	-
median	11.0	11.5	12.6	13.8	15.0	15.6	16.6	18.0	18.0	-
10th	-	9.7	11.2	12.0	12.8	14.5	15.3	15.5	-	-
testosterone (ng/dl)										
N	4	66	104	76	69	59	32	5	0	0
90th	-	160	336	430	517	575	787	-	-	-
median	36	36	86	148	259	400	481	569	-	-
10th	-	13	21	38	44	110	327	-	-	-
DHEAS (ng/ml)										
N	0	4	67	43	32	31	30	6	0	0
90th	-	-	2320	3160	3200	3560	5000	-	-	-
median	-	464	1160	1480	1650	1840	1940	2270	-	-
10th	-	-	420	720	560	1040	900	-	-	-

Entries are number of measurements and percentiles. Only the median is shown if $n \leq 10$.

Table 4: Mean anthropometric and pubertal measures by p,p'-DDE in specific age ranges

p,p'-DDE (μ g/g lipid)	<14 years	14-<17 years	17-20 years
height (cm)			
<3	151 \pm 1.8 (28)	168 \pm 1.9 (26)	174 \pm 1.5 (25)
3-<6	154 \pm 0.9 (76)	168 \pm 0.9 (86)	176 \pm 0.8 (67)
6-<9	155 \pm 1.2 (50)	170 \pm 1.3 (50)	176 \pm 1.0 (51)
9-<12	157 \pm 2.0 (27)	173 \pm 1.7 (24)	177 \pm 1.5 (24)
12+	152 \pm 1.9 (18)	168 \pm 2.1 (21)	173 \pm 1.4 (19)
height ratio (%)			
<3	50.8 \pm 0.3 (28)	51.1 \pm 0.4 (26)	51.2 \pm 0.3 (25)
3-<6	50.8 \pm 0.1 (76)	50.6 \pm 0.2 (84)	51.0 \pm 0.2 (66)
6-<9	50.7 \pm 0.2 (50)	50.3 \pm 0.2 (50)	50.7 \pm 0.2 (51)
9-<12	50.2 \pm 0.3 (27)	50.3 \pm 0.3 (24)	50.3 \pm 0.3 (24)
12+	50.7 \pm 0.3 (18)	50.2 \pm 0.3 (21)	50.9 \pm 0.4 (18)
BMI (kg/m ²)			
<3	17.9 \pm 0.5 (28)	21.6 \pm 1.0 (26)	21.5 \pm 0.6 (25)
3-<6	20.0 \pm 0.5 (76)	20.8 \pm 0.5 (85)	22.9 \pm 0.4 (67)
6-<9	18.9 \pm 0.4 (50)	20.4 \pm 0.5 (50)	22.6 \pm 0.4 (51)
9-<12	18.3 \pm 0.5 (27)	21.3 \pm 1.0 (24)	22.3 \pm 0.9 (24)
12+	18.2 \pm 0.5 (18)	19.7 \pm 0.6 (21)	21.2 \pm 0.6 (19)
triceps skinfold thickness (mm)			
<3	8.5 \pm 0.6 (28)	10.3 \pm 1.5 (26)	7.5 \pm 0.6 (25)
3-<6	11.0 \pm 0.8 (76)	8.4 \pm 0.5 (86)	8.9 \pm 0.6 (67)
6-<9	9.4 \pm 0.7 (50)	8.5 \pm 0.7 (50)	8.7 \pm 0.6 (51)
9-<12	8.9 \pm 0.9 (27)	9.1 \pm 1.5 (24)	8.6 \pm 1.4 (24)
12+	8.1 \pm 0.7 (18)	8.2 \pm 0.8 (21)	7.2 \pm 0.8 (19)

central adiposity (%)			
<3	46.2 ± 0.8 (28)	49.1 ± 1.0 (26)	56.8 ± 0.9 (25)
3-<6	47.6 ± 0.6 (75)	51.9 ± 0.6 (86)	56.1 ± 0.6 (67)
6-<9	47.7 ± 0.7 (50)	51.2 ± 0.7 (50)	55.3 ± 0.8 (51)
9-<12	48.0 ± 0.7 (27)	50.8 ± 1.0 (24)	58.1 ± 0.9 (24)
12+	46.5 ± 1.1 (18)	50.7 ± 1.0 (21)	57.2 ± 1.3 (19)
skeletal age (yrs)			
<3	12.4 ± 0.3 (28)	15.8 ± 0.3 (24)	
3-<6	13.1 ± 0.1 (72)	15.4 ± 0.1 (73)	
6-<9	13.0 ± 0.2 (50)	15.4 ± 0.2 (43)	
9-<12	13.1 ± 0.3 (27)	15.9 ± 0.2 (22)	
12+	13.0 ± 0.3 (18)	15.6 ± 0.3 (21)	
testosterone (ng/dl)			
<3	124 ± 26 (26)	376 ± 33 (18)	
3-<6	136 ± 17 (63)	321 ± 28 (43)	
6-<9	172 ± 25 (42)	346 ± 46 (25)	
9-<12	111 ± 19 (24)	491 ± 103 (11)	
12+	142 ± 41 (13)	333 ± 52 (15)	
DHEAS (ng/ml)			
<3	1355 ± 184 (20)	2248 ± 274 (16)	
3-<6	1431 ± 142 (40)	1901 ± 168 (35)	
6-<9	1428 ± 186 (28)	1911 ± 218 (19)	
9-<12	1236 ± 186 (18)	2352 ± 476 (10)	
12+	1240 ± 180 (8)	2186 ± 492 (13)	

For each boy, all available measurements in the specified age range are averaged. Entries are mean ± standard error of these averages (number of boys). Skeletal age and hormone concentrations are not shown in the upper age range since they were available for few boys.

Table 5: Regression of anthropometric and pubertal measures on maternal p,p'-DDE and other predictors

predictor	category /units	outcome							
		height (cm)	height ratio (%)	BMI (kg/m ²)	triceps (log mm)	central adiposity (%)	skeletal age (yrs)	testosterone (log ng/dl)	DHEAS (ng/ml)
maternal	<3	0 (ref)	0 (ref)	0 (ref)	0 (ref)	0 (ref)	0 (ref)	0 (ref)	0 (ref)
p,p'-DDE	3-<6	1.1 ± 1.3	0.1 ± 0.2	0.9 ± 0.7	0.06 ± 0.08	1.1 ± 0.9	0.4 ± 0.2	0.0 ± 0.2	-158 ± 217
(□g/g lipid)	6-<9	1.0 ± 1.5	0.1 ± 0.2	0.2 ± 0.8	0.00 ± 0.09	0.5 ± 1.0	0.3 ± 0.3	0.1 ± 0.2	-140 ± 252
	9-<12	2.2 ± 1.7	0.0 ± 0.3	0.6 ± 0.9	0.00 ± 0.10	1.7 ± 1.2	0.5 ± 0.3	0.0 ± 0.2	-109 ± 281
	12+	0.4 ± 1.8	0.0 ± 0.3	-0.4 ± 0.9	-0.01 ± 0.11	0.6 ± 1.2	0.2 ± 0.3	0.1 ± 0.2	-148 ± 307
maternal height	per 10 cm	4.4 ± 0.7**	-0.5 ± 0.1**	-0.2 ± 0.4	0.05 ± 0.04	0.0 ± 0.5	-0.1 ± 0.1	-0.1 ± 0.1	-32 ± 114
maternal BMI	per 10 kg/m ²	2.0 ± 1.0*	0.0 ± 0.2	2.2 ± 0.5**	0.25 ± 0.06**	-0.3 ± 0.7	0.2 ± 0.2	0.2 ± 0.1	280 ± 179
race	white	-0.9 ± 1.3	2.1 ± 0.2	0.4 ± 0.7	0.15 ± 0.08	-2.8 ± 0.9	0.0 ± 0.2	0.0 ± 0.2	-110 ± 227
	black	0 (ref)	0 (ref)**	0 (ref)	0 (ref)*	0 (ref)**	0 (ref)	0 (ref)	0 (ref)
# older siblings	0	0 (ref)**	0 (ref)*	0 (ref)	0 (ref)**	0 (ref)	0 (ref)	0 (ref)	0 (ref)
	1	-1.1 ± 1.3	0.0 ± 0.2	-0.6 ± 0.7	-0.11 ± 0.08	1.4 ± 0.9	-0.3 ± 0.2	-0.2 ± 0.2	-320 ± 235
	2+	-3.2 ± 1.4	0.4 ± 0.2	-0.9 ± 0.7	-0.21 ± 0.08	1.7 ± 0.9	-0.5 ± 0.2	-0.2 ± 0.2	-522 ± 245

maternal smoking no		0 (ref)	0 (ref)	0 (ref)	0 (ref)	0 (ref)	0 (ref)	0 (ref)	0 (ref)
in pregnancy	yes	-0.2 ± 0.8	-0.2 ± 0.1	0.3 ± 0.5	0.02 ± 0.05	0.7 ± 0.6	0.0 ± 0.1	0.0 ± 0.1	60 ± 152
breast-fed	no	0.4 ± 1.7	-0.3 ± 0.3	0.5 ± 0.9	0.12 ± 0.10	-0.7 ± 1.2	-0.1 ± 0.3	0.1 ± 0.2	-247 ± 285
	yes	0 (ref)	0 (ref)	0 (ref)	0 (ref)	0 (ref)	0 (ref)	0 (ref)	0 (ref)
family SEI	5.1+	2.4 ± 1.4	0.4 ± 0.2	1.1 ± 0.7	0.12 ± 0.08	-0.3 ± 0.9	0.4 ± 0.2	0.4 ± 0.2	207 ± 238
	2.6-5.0	2.5 ± 1.2	0.1 ± 0.2	0.7 ± 0.6	0.04 ± 0.07	0.0 ± 0.8	0.4 ± 0.2	0.2 ± 0.2	184 ± 204
	0-2.5	0 (ref)*	0 (ref)	0 (ref)	0 (ref)	0 (ref)	0 (ref)	0 (ref)*	0 (ref)
maternal age at	13-19	-0.9 ± 1.8	-0.1 ± 0.3	-0.7 ± 1.0	-0.20 ± 0.11	1.7 ± 1.2	-0.4 ± 0.3	0.2 ± 0.2	-340 ± 318
enrollment (yrs)	20-24	-0.6 ± 1.3	-0.3 ± 0.2	-0.5 ± 0.7	-0.08 ± 0.08	0.4 ± 0.9	-0.2 ± 0.2	-0.1 ± 0.2	-122 ± 237
	25-29	0.2 ± 1.4	-0.4 ± 0.2	0.2 ± 0.7	-0.05 ± 0.08	0.5 ± 1.0	0.0 ± 0.2	0.2 ± 0.2	-252 ± 243
	30+	0 (ref)	0 (ref)	0 (ref)	0 (ref)	0 (ref)	0 (ref)	0 (ref)	0 (ref)
maternal age at	8-11	0 (ref)	0 (ref)*	0 (ref)	0 (ref)	0 (ref)	0 (ref)	0 (ref)*	0 (ref)
menarche (yrs)	12	-0.6 ± 1.2	-0.5 ± 0.2	0.0 ± 0.6	0.03 ± 0.07	-1.2 ± 0.8	-0.3 ± 0.2	-0.5 ± 0.2	-77 ± 212
	13	0.3 ± 1.3	-0.4 ± 0.2	-0.4 ± 0.7	0.00 ± 0.08	-0.9 ± 0.9	-0.2 ± 0.2	-0.3 ± 0.2	47 ± 228
	14+	1.0 ± 1.3	-0.4 ± 0.2	-0.3 ± 0.7	0.00 ± 0.08	-1.3 ± 0.9	-0.4 ± 0.2	-0.3 ± 0.2	-52 ± 223

Entries are regression coefficients (\pm standard error), adjusted for all others shown and also for age of boy at measurement and correlation among multiple measures of the same boy. * $p < .10$, ** $p < .05$, for tests of whether discrete groups differ or continuous slope is nonzero.